

Dr A.L.Sabin,
Children's Hospital
Cincinnati 29
Ohio

December 25, 1961
Belgrade

Dear Dr. Sabin:

Few days ago I came back from the United States, where I went to attend the meeting on measles immunization in Bethesda. The meeting was held 7-10 November, and afterwards I spent some time in N.I.H., then went to ~~xxx~~ Boston and New-York. On my way home I visited London too. This is the reason why I answer so late your letter of November 8. I am so sorry I could not give you clinical data on the reported cases, because the time does not permit me, what I am going to do later, as soon as possible.

You will find enclosed here some data I had with me, and I hope it may be of help for you.

As far as I know polio morbidity rate for the whole of Yugoslavia was about the same as the one in Serbia. The whole country was vaccinated with live polio vaccine, ^{Sabin type} except Croatia where Koprowsky type 1 and type 3 were used, type 2 was not used in Croatia at all.

As to the laboratory data from other republics than Serbia-I do not know anything. It has been planned-all reported cases to be tested virologically in Serbia, but unfortunately I was not able to obtain the specimens from all reported cases.

As soon as I get the data you want-I will write you.

With the best wishes in New Year

sincerely yours,

M.V. Milovanovic

Dr. M.V. Milovanović,
Institute of Hygiene
Belgrade 7
Yugoslavia

See Stojkovic letter of Sept. 9 —

"In Serbia after vaccination
not a single proved
case" — reported cases
many clinically, not etc.
4.5 million polio.

Vaccination in 1961 — according to Stojkovic

Type 1 — began Feb. 7

" 3 — end of March and early April

" 2 — Middle May

See p. 145

In Bosnia Diph Tet. & Pertussis
given same day.

in Lyon Proceedings

month	No reported vaccinated nonvacc. totals			No. tested vaccinated nonvaccin. totals		
I	0	7	7	0	2 /type I- I negative- I/	2
2	2	4	6	2 /type I-2/	2 /type I-I neg.- I/	4
3	I	4	5	I /neg.- I/	I /neg.-I/	2
4	I	I	2	0	I /neg.-I/	I
5	2	0	2	I /neg.-I/	0	I
6	I	3	4	I /neg.-I/	I /non-polio-I/	2
7	3	I	4	2 /neg.-2/	I /neg.-I/	3
8	3	0	3	I /neg.-I/	0	I
9	0	2	2	0	I /type 2-I/	I
IO	I	0	I	I /neg.-I/	0	I
II	I	2	3	I /neg.-I/	2 /neg.-2/	3

* Both cases were fed type I two days before the onset of the disease.

** Fatal case where the nonpolio agent was isolated from the spinal cord.

March to Nov. 13
July - Nov 8
Vaccin. Started with Type 1
on Feb. 7

13

0/5
+3

1/4 - Type 2
+2
8

	No. reported	No. tested	No. Polio Recovered
Jan. - before vacc.	7	2	1 (Type 1)
Feb. During Vacc.	6	4	3 (Type 1) - 2/2 Type 1 from cases that received Type 1 before onset Type 1
March Apr May	9	4	0
June, July, Aug	11	6	1 ← Non Polio
Sept Oct Nov.	6	5	1 ← Type 2